

Phonon

Mitsuaki Kawamura

September 14, 2022

1 Single harmonic Oscillator

The hamiltonian operator is

$$\hat{H} = \frac{\hat{p}^2}{2M} + \frac{C\hat{u}^2}{2}, \quad (1)$$

where \hat{u} and \hat{p} are the position and the momentum operator. These operators satisfies the following commutation relation:

$$[\hat{u}, \hat{p}] \equiv \hat{u}\hat{p} - \hat{p}\hat{u} = i \quad (2)$$

Then we introduce the creation- and anihilation- operator (\hat{b}^\dagger and \hat{b}) as follows

$$\hat{u} \equiv \frac{1}{(4MC)^{1/4}}(\hat{b} + \hat{b}^\dagger), \quad (3)$$

$$\hat{p} \equiv \left(\frac{MC}{4}\right)^{1/4}(-i\hat{b} + i\hat{b}^\dagger). \quad (4)$$

The commutation relation of \hat{b}^\dagger and \hat{b}

$$[\hat{b}, \hat{b}] = 0, [\hat{b}, \hat{b}^\dagger] = 1 \quad (5)$$

lead to the original commutation relation of \hat{u} and \hat{p} as

$$[\hat{u}, \hat{p}] = \frac{i}{2}(-[\hat{b}, \hat{b}] + [\hat{b}, \hat{b}^\dagger] - [\hat{b}^\dagger, \hat{b}] + [\hat{b}^\dagger, \hat{b}^\dagger]) = i, \quad (6)$$

and the hamiltonian becomes

$$\begin{aligned} \hat{H} &= \frac{1}{4} \left(\frac{(MC)^{1/2}}{M} (-i\hat{b} + i\hat{b}^\dagger)(-i\hat{b} + i\hat{b}^\dagger) + \frac{C}{(MC)^{1/2}} (\hat{b} + \hat{b}^\dagger)(\hat{b} + \hat{b}^\dagger) \right) \\ &= \frac{1}{2} \left(\frac{C}{M} \right)^{1/2} (\hat{b}\hat{b}^\dagger + \hat{b}^\dagger\hat{b}) = \omega \left(\hat{b}^\dagger\hat{b} + \frac{1}{2} \right), \end{aligned} \quad (7)$$

where $\omega \equiv (C/M)^{1/2}$

2 Coupled Harmonic Oscillator

The hamiltonian operator is

$$\hat{H} = \frac{1}{2} \sum_s \frac{\hat{p}_s^2}{M_s} + \frac{1}{2} \sum_{ss'} \hat{u}_s C_{ss'} \hat{u}_{s'}, \quad (8)$$

where \hat{u}_s and \hat{p}_s are the position and the momentum operator. These operators satisfies the following commutation relation:

$$[\hat{u}_s, \hat{p}_{s'}] \equiv i\delta_{ss'}. \quad (9)$$

Then, we introduce the creation- and anihilation- operator (\hat{b}_ν^\dagger and \hat{b}_ν) as follows

$$\hat{u}_s \equiv \sum_\nu \frac{v_{s\nu}}{(M_s \omega_\nu)^{1/2}} (\hat{b}_\nu + \hat{b}_\nu^\dagger), \quad (10)$$

$$\hat{p}_s \equiv \sum_\nu (M_s \omega_\nu)^{1/2} v_{s\nu} (-i\hat{b}_\nu + i\hat{b}_\nu^\dagger), \quad (11)$$

where $v_{s\nu}$ and ω_ν^2 are eigenvectors and eigenvalues of rescaled force constant as

$$\sum_{s'} \frac{C_{ss'}}{(M_s M_{s'})^{1/2}} v_{s'\nu} = \omega_\nu^2 v_{s\nu} \quad (12)$$

The commutation relation of \hat{b}_ν^\dagger and \hat{b}_ν

$$[\hat{b}_\nu, \hat{b}_{\nu'}] = 0, [\hat{b}_\nu, \hat{b}_{\nu'}^\dagger] = \delta_{\nu\nu'} \quad (13)$$

lead to the original commutation relation of \hat{u}_s and \hat{p}_s as

$$[\hat{u}_s, \hat{p}_{s'}] = \sum_{\nu\nu'} v_{s\nu} v_{s'\nu'} \frac{i}{2} (-[\hat{b}_\nu, \hat{b}_{\nu'}] + [\hat{b}_\nu, \hat{b}_{\nu'}^\dagger] - [\hat{b}_\nu^\dagger, \hat{b}_{\nu'}] + [\hat{b}_\nu^\dagger, \hat{b}_{\nu'}^\dagger]) = i \sum_\nu v_{s\nu} v_{s'\nu} = i\delta_{ss'}, \quad (14)$$

and the hamiltonian becomes

$$\begin{aligned} \hat{H} &= \frac{1}{2} \sum_{\nu\nu'} (\omega_\nu \omega_{\nu'})^{1/2} (-i\hat{b}_\nu + i\hat{b}_\nu^\dagger) (-i\hat{b}_{\nu'} + i\hat{b}_{\nu'}^\dagger) \sum_s \frac{M_s}{M_s} v_{s\nu} v_{s\nu'} \\ &+ \frac{1}{2} \sum_{\nu\nu'} (\hat{b}_\nu + \hat{b}_\nu^\dagger) (\hat{b}_{\nu'} + \hat{b}_{\nu'}^\dagger) \frac{1}{(\omega_\nu \omega_{\nu'})^{1/2}} \sum_{ss'} v_{s\nu} \frac{C_{ss'}}{(M_s M_{s'})^{1/2}} v_{s'\nu'} \\ &= \frac{1}{2} \sum_{\nu\nu'} (\omega_\nu \omega_{\nu'})^{1/2} (-\hat{b}_\nu \hat{b}_{\nu'} + \hat{b}_\nu \hat{b}_{\nu'}^\dagger + \hat{b}_\nu^\dagger \hat{b}_{\nu'} - \hat{b}_\nu^\dagger \hat{b}_{\nu'}^\dagger) \sum_s v_{s\nu} v_{s\nu'} \\ &+ \frac{1}{2} \sum_{\nu\nu'} (\hat{b}_\nu \hat{b}_{\nu'} + \hat{b}_\nu \hat{b}_{\nu'}^\dagger + \hat{b}_\nu^\dagger \hat{b}_{\nu'} + \hat{b}_\nu^\dagger \hat{b}_{\nu'}^\dagger) \frac{\omega_{\nu'}^2}{(\omega_\nu \omega_{\nu'})^{1/2}} \sum_s v_{s\nu} v_{s\nu'} \\ &= \frac{1}{2} \sum_\nu \omega_\nu (\hat{b}_\nu \hat{b}_\nu^\dagger + \hat{b}_\nu^\dagger \hat{b}_\nu) = \sum_\nu \omega_\nu \left(\hat{b}_\nu^\dagger \hat{b}_\nu + \frac{1}{2} \right), \end{aligned} \quad (15)$$

3 Periodic Coupled Harmonic Oscillator

The hamiltonian operator is

$$\hat{H} = \frac{1}{2} \sum_{\mathbf{T}s\alpha} \frac{\hat{p}_{\mathbf{T}s\alpha}^2}{M_s} + \frac{1}{2} \sum_{\mathbf{T}s\alpha \mathbf{T}'s'\alpha'} \hat{u}_{\mathbf{T}s\alpha} C_{\mathbf{T}s\alpha \mathbf{T}'s'\alpha'} \hat{u}_{\mathbf{T}'s'\alpha'}, \quad (16)$$

where $\hat{u}_{\mathbf{T}s\alpha}$ and $\hat{p}_{\mathbf{T}s\alpha}$ are the position and the momentum operator. These operators satisfies the following commutation relation:

$$[\hat{u}_{\mathbf{T}s\alpha}, \hat{p}_{\mathbf{T}'s'\alpha'}] \equiv i\delta_{\mathbf{T}\mathbf{T}'} \delta_{ss'} \delta_{\alpha\alpha'}. \quad (17)$$

The Fourier-transformed operators are defined as follows:

$$\hat{U}_{\mathbf{q}s\alpha} \equiv \frac{1}{N_C^{1/2}} \sum_{\mathbf{T}} \hat{u}_{\mathbf{T}s\alpha} e^{i\mathbf{q}\cdot\mathbf{T}}, \hat{P}_{\mathbf{q}s\alpha} \equiv \frac{1}{N_C^{1/2}} \sum_{\mathbf{T}} \hat{p}_{\mathbf{T}s\alpha} e^{i\mathbf{q}\cdot\mathbf{T}} \quad (18)$$

$$\hat{u}_{\mathbf{T}s\alpha} = \frac{1}{N_C^{1/2}} \sum_{\mathbf{q}} \hat{U}_{\mathbf{q}s\alpha} e^{-i\mathbf{q}\cdot\mathbf{T}}, \hat{p}_{\mathbf{T}s\alpha} = \frac{1}{N_C^{1/2}} \sum_{\mathbf{q}} \hat{P}_{\mathbf{q}s\alpha} e^{-i\mathbf{q}\cdot\mathbf{T}}, \quad (19)$$

where N_C is the number of cells within the Born—von Karman boundary condition. They also satisfy the commutation relation.

$$[\hat{U}_{\mathbf{q}s\alpha}, \hat{P}_{\mathbf{q}'s'\alpha'}^\dagger] = \frac{1}{N_C} \sum_{\mathbf{T}\mathbf{T}'} e^{i\mathbf{q}\cdot\mathbf{T}} e^{-i\mathbf{q}'\cdot\mathbf{T}'} [\hat{u}_{\mathbf{T}s\alpha}, \hat{p}_{\mathbf{T}'s'\alpha'}] = i \frac{1}{N_C} \sum_{\mathbf{T}} e^{i(\mathbf{q}-\mathbf{q}')\cdot\mathbf{T}} \delta_{ss'} \delta_{\alpha\alpha'} \quad (20)$$

$$= i \delta_{\mathbf{q}\mathbf{q}'} \delta_{ss'} \delta_{\alpha\alpha'}. \quad (21)$$

The hamiltonian becomes

$$\begin{aligned} \hat{H} &= \frac{1}{2N_C} \sum_{\mathbf{q}\mathbf{q}'\mathbf{T}s\alpha} \frac{\hat{P}_{\mathbf{q}s\alpha} \hat{P}_{\mathbf{q}'s'\alpha'} e^{i(\mathbf{q}+\mathbf{q}')\cdot\mathbf{T}}}{M_s} + \frac{1}{2N_C} \sum_{\mathbf{q}\mathbf{q}'s\alpha s'\alpha'} \hat{U}_{\mathbf{q}s\alpha} \hat{U}_{\mathbf{q}'s'\alpha'} \sum_{\mathbf{T}\mathbf{T}'} C_{\mathbf{T}s\alpha\mathbf{T}'s'\alpha'} e^{i\mathbf{q}\cdot\mathbf{T}} e^{i\mathbf{q}'\cdot\mathbf{T}'} \\ &= \frac{1}{2} \sum_{\mathbf{q}\mathbf{q}'\mathbf{T}s\alpha} \frac{\hat{P}_{\mathbf{q}s\alpha}^\dagger \hat{P}_{\mathbf{q}s\alpha}}{M_s} + \frac{1}{2N_C} \sum_{\mathbf{q}\mathbf{q}'s\alpha s'\alpha'} \hat{U}_{\mathbf{q}s\alpha} \hat{U}_{\mathbf{q}'s'\alpha'} \sum_{\mathbf{T}\mathbf{T}'} C_{0s\alpha(\mathbf{T}'-\mathbf{T})s'\alpha'} e^{i\mathbf{q}'\cdot(\mathbf{T}'-\mathbf{T})} e^{i(\mathbf{q}+\mathbf{q}')\cdot\mathbf{T}} \\ &= \sum_{\mathbf{q}} \left(\frac{1}{2} \sum_{s\alpha} \frac{\hat{P}_{\mathbf{q}s\alpha}^\dagger \hat{P}_{\mathbf{q}s\alpha}}{M_s} + \frac{1}{2} \sum_{s\alpha s'\alpha'} \hat{U}_{\mathbf{q}s\alpha}^\dagger \tilde{C}_{\mathbf{q}s\alpha s'\alpha'} \hat{U}_{\mathbf{q}s'\alpha'} \right), \end{aligned} \quad (22)$$

where

$$\tilde{C}_{\mathbf{q}s\alpha s'\alpha'} \equiv \sum_{\mathbf{T}} C_{0s\alpha\mathbf{T}s'\alpha'} e^{i\mathbf{q}\cdot\mathbf{T}} \quad (23)$$

With the same discussion in the previous section, we obtain the following results:

$$\hat{U}_{\mathbf{q}s\alpha} \equiv \sum_{\nu} \frac{v_{s\alpha\mathbf{q}\nu}}{(M_s \omega_{\mathbf{q}\nu})^{1/2}} (\hat{b}_{\mathbf{q}\nu} + \hat{b}_{\mathbf{q}\nu}^\dagger), \quad (24)$$

$$\hat{P}_{\mathbf{q}s\alpha} \equiv \sum_{\nu} (M_s \omega_{\mathbf{q}\nu})^{1/2} v_{s\alpha\mathbf{q}\nu} (-i \hat{b}_{\mathbf{q}\nu} + i \hat{b}_{\mathbf{q}\nu}^\dagger), \quad (25)$$

$$\sum_{s'\alpha'} \frac{\tilde{C}_{\mathbf{q}s\alpha s'\alpha'}}{(M_s M_{s'})^{1/2}} v_{s'\alpha'\mathbf{q}\nu} = \omega_{\mathbf{q}\nu}^2 v_{s\alpha\mathbf{q}\nu} \quad (26)$$

$$\hat{H} = \sum_{\mathbf{q}\nu} \omega_{\mathbf{q}\nu} \left(\hat{b}_{\mathbf{q}\nu}^\dagger \hat{b}_{\mathbf{q}\nu} + \frac{1}{2} \right) \quad (27)$$

$$\begin{aligned} [\hat{U}_{\mathbf{q}s\alpha}, \hat{P}_{\mathbf{q}'s'\alpha'}^\dagger] &= \sum_{\nu\nu'} v_{s\alpha\mathbf{q}\nu} v_{s'\alpha'\mathbf{q}'\nu'}^* \frac{i}{2} (-[\hat{b}_{\mathbf{q}\nu}, \hat{b}_{\mathbf{q}'\nu'}] + [\hat{b}_{\mathbf{q}\nu}, \hat{b}_{\mathbf{q}'\nu'}^\dagger] - [\hat{b}_{\mathbf{q}\nu}^\dagger, \hat{b}_{\mathbf{q}'\nu'}] + [\hat{b}_{\mathbf{q}\nu}^\dagger, \hat{b}_{\mathbf{q}'\nu'}^\dagger]) \\ &= i \delta_{\mathbf{q}\mathbf{q}'} \sum_{\nu} v_{s\alpha\mathbf{q}\nu} v_{s'\alpha'\mathbf{q}'\nu}^* = i \delta_{\mathbf{q}\mathbf{q}'} \delta_{ss'} \delta_{\alpha\alpha'}, \end{aligned} \quad (28)$$

4 Electron-phonon vertex

Electron-nuclear Hamiltonian in 2nd quantization representation

$$\hat{H}_{en} = \sum_{\sigma} \int d^3r \hat{\psi}_{\sigma}(\mathbf{r})^\dagger \hat{\psi}_{\sigma}(\mathbf{r}) V(\mathbf{r}; \{\hat{R}_{\mathbf{T}s\alpha}\}), \quad (29)$$

where $\hat{R}_{\mathbf{T}s\alpha}$ is the position operator of nuclear as

$$\hat{R}_{\mathbf{T}s\alpha} = R_{\mathbf{T}s\alpha}^0 + \hat{u}_{\mathbf{T}s\alpha}, \quad (30)$$

$R_{\mathbf{T}s\alpha}^0$ is equilibrium position.

We expand $V(\mathbf{r}; \{\hat{R}_{\mathbf{T}s\alpha}\})$ around $R_{\mathbf{T}s\alpha}^0$ and obtain

$$\hat{H}_{en} = \sum_{\sigma} \int d^3r \hat{\psi}_{\sigma}(\mathbf{r})^{\dagger} \hat{\psi}_{\sigma}(\mathbf{r}) V(\mathbf{r}; \{R_{\mathbf{T}s\alpha}^0\}) + \sum_{\sigma} \int d^3r \hat{\psi}_{\sigma}(\mathbf{r})^{\dagger} \hat{\psi}_{\sigma}(\mathbf{r}) \sum_{\mathbf{T}s\alpha} \frac{\partial V(\mathbf{r}; \{R^0\})}{\partial R_{\mathbf{T}s\alpha}^0} \hat{u}_{\mathbf{T}s\alpha} + O(u^2). \quad (31)$$

The first term is the interaction between electron and fixed nuclear, and the second term is the electron-phonon interaction \hat{H}_{ep} . By expanding $\hat{\psi}_{\sigma}$ with Bloch orbitals $\varphi_{n\mathbf{k}}$

$$\hat{\psi}_{\sigma}(\mathbf{r}) = \sum_{n\mathbf{k}} \varphi_{n\mathbf{k}}(\mathbf{r}) \hat{c}_{n\mathbf{k}\sigma}, \quad (32)$$

we obtain

$$\hat{H}_{ep} = \sum_{\sigma nn'\mathbf{k}\mathbf{k}'\mathbf{T}s\alpha} \hat{c}_{n\mathbf{k}\sigma}^{\dagger} \hat{c}_{n'\mathbf{k}'\sigma} \hat{u}_{\mathbf{T}s\alpha} \int d^3r \varphi_{n\mathbf{k}}^*(\mathbf{r}) \varphi_{n'\mathbf{k}'}(\mathbf{r}) \frac{\partial V(\mathbf{r}; \{R^0\})}{\partial R_{\mathbf{T}s\alpha}^0}. \quad (33)$$

By using Eqs (19, 24) and $\varphi_{n\mathbf{k}}(\mathbf{r}) \equiv N_C^{-1/2} e^{i\mathbf{k}\cdot\mathbf{r}} \chi_{n\mathbf{k}}(\mathbf{r})$, we obtain

$$\hat{H}_{ep} = \sum_{\sigma nn'\mathbf{k}\mathbf{k}'\mathbf{q}\nu} \hat{c}_{n\mathbf{k}\sigma}^{\dagger} \hat{c}_{n'\mathbf{k}'\sigma} (\hat{b}_{\mathbf{q}\nu} + \hat{b}_{\mathbf{q}\nu}^{\dagger}) g_{n\mathbf{k}n'\mathbf{k}'}^{\mathbf{q}\nu}, \quad (34)$$

where $g_{n\mathbf{k}n'\mathbf{k}'}^{\mathbf{q}\nu}$ is the electron-phonon vertex as

$$\begin{aligned} g_{n\mathbf{k}n'\mathbf{k}'}^{\mathbf{q}\nu} &\equiv \sum_{\mathbf{T}s\alpha} e^{-i\mathbf{q}\cdot\mathbf{T}} \frac{v_{s\alpha\mathbf{q}\nu}}{(N_C^3 M_s \omega_{\mathbf{q}\nu})^{1/2}} \int d^3r e^{i(\mathbf{k}'-\mathbf{k})\cdot\mathbf{r}} \chi_{n\mathbf{k}}^*(\mathbf{r}) \chi_{n'\mathbf{k}'}(\mathbf{r}) \frac{\partial V(\mathbf{r}; \{R^0\})}{\partial R_{\mathbf{T}s\alpha}^0} \\ &= \sum_{\mathbf{T}\mathbf{T}'s\alpha} e^{-i\mathbf{q}\cdot\mathbf{T}} \frac{v_{s\alpha\mathbf{q}\nu}}{(N_C^3 M_s \omega_{\mathbf{q}\nu})^{1/2}} \int_{\text{cell}} d^3r e^{i(\mathbf{k}'-\mathbf{k})\cdot(\mathbf{r}+\mathbf{T}')} \chi_{n\mathbf{k}}^*(\mathbf{r}) \chi_{n'\mathbf{k}'}(\mathbf{r}) \frac{\partial V(\mathbf{r}+\mathbf{T}'; \{R^0\})}{\partial R_{\mathbf{T}s\alpha}^0} \\ &= \sum_{\mathbf{T}} e^{i(\mathbf{k}'-\mathbf{k}-\mathbf{q})\cdot\mathbf{T}} \sum_{\mathbf{T}'s\alpha} \frac{v_{s\alpha\mathbf{q}\nu}}{(N_C^3 M_s \omega_{\mathbf{q}\nu})^{1/2}} \int_{\text{cell}} d^3r e^{i(\mathbf{k}'-\mathbf{k})\cdot(\mathbf{r}+\mathbf{T}'-\mathbf{T})} \chi_{n\mathbf{k}}^*(\mathbf{r}) \chi_{n'\mathbf{k}'}(\mathbf{r}) \frac{\partial V(\mathbf{r}; \{R^0\})}{\partial R_{(\mathbf{T}-\mathbf{T}')s\alpha}^0} \\ &= \delta_{\mathbf{k}',\mathbf{k}+\mathbf{q}} \sum_{s\alpha} \frac{v_{s\alpha\mathbf{q}\nu}}{(N_C M_s \omega_{\mathbf{q}\nu})^{1/2}} \int_{\text{cell}} d^3r \chi_{n\mathbf{k}}^*(\mathbf{r}) \chi_{n'\mathbf{k}+\mathbf{q}}(\mathbf{r}) \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{r}-\mathbf{T})} \frac{\partial V(\mathbf{r}; \{R^0\})}{\partial R_{\mathbf{T}s\alpha}^0}. \end{aligned} \quad (35)$$

If we use V_{KS} alternative to V , we obtain the screened electron-phonon vertex. The screened deformation potential $\sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{r}-\mathbf{T})} \partial V_{\text{KS}}(\mathbf{r}; \{R^0\}) / \partial R_{\mathbf{T}s\alpha}^0$ is obtained as a biproduct of DFPT calculation.

This deformation potential has lattice periodicity as

$$\sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{r}+\mathbf{T}'-\mathbf{T})} \frac{\partial V(\mathbf{r}+\mathbf{T}'; \{R^0\})}{\partial R_{\mathbf{T}s\alpha}^0} = \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{r}-\mathbf{T})} \frac{\partial V(\mathbf{r}; \{R^0\})}{\partial R_{\mathbf{T}s\alpha}^0} \quad (36)$$

5 Density functional perturbation theory for lattice

Force constant

$$\begin{aligned} C_{\mathbf{T}s\alpha\mathbf{T}'s'\alpha'} &\equiv \frac{\partial^2 E}{\partial R_{\mathbf{T}s\alpha}^0 \partial R_{\mathbf{T}'s'\alpha'}^0} = -\frac{\partial F_{\mathbf{T}s\alpha}}{\partial R_{\mathbf{T}'s'\alpha'}^0} = \frac{\partial}{\partial R_{\mathbf{T}'s'\alpha'}^0} \left(-F_{\mathbf{T}s\alpha}^{\text{C}} + \int d^3r \rho(\mathbf{r}) \frac{\partial V(\mathbf{r}; \{R^0\})}{\partial R_{\mathbf{T}s\alpha}^0} \right) \\ &= \frac{\partial^2 E_{\text{C}}}{\partial R_{\mathbf{T}s\alpha}^0 \partial R_{\mathbf{T}'s'\alpha'}^0} + \int d^3r \rho(\mathbf{r}) \frac{\partial^2 V(\mathbf{r}; \{R^0\})}{\partial R_{\mathbf{T}s\alpha}^0 \partial R_{\mathbf{T}'s'\alpha'}^0} + \int d^3r \frac{\partial \rho(\mathbf{r})}{\partial R_{\mathbf{T}'s'\alpha'}^0} \frac{\partial V(\mathbf{r}; \{R^0\})}{\partial R_{\mathbf{T}s\alpha}^0} \end{aligned} \quad (37)$$

Dynamical matrix

$$\begin{aligned}
\tilde{C}_{\mathbf{q}s\alpha s'\alpha'} &\equiv \sum_{\mathbf{T}} C_{\mathbf{0}s\alpha\mathbf{T}s'\alpha'} e^{i\mathbf{q}\cdot\mathbf{T}} \\
&= \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot\mathbf{T}} \left(\frac{\partial^2 E_C}{\partial R_{s\alpha}^0 \partial R_{\mathbf{T}s'\alpha'}^0} + \int d^3r \rho(\mathbf{r}) \frac{\partial^2 V(\mathbf{r}; \{R^0\})}{\partial R_{\mathbf{0}s\alpha}^0 \partial R_{\mathbf{T}s'\alpha'}^0} \right) + \int d^3r \frac{\partial V(\mathbf{r}; \{R^0\})}{\partial R_{\mathbf{0}s\alpha}^0} \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot\mathbf{T}} \frac{\partial \rho(\mathbf{r})}{\partial R_{\mathbf{T}s'\alpha'}^0}
\end{aligned} \tag{38}$$

Monochromatic perturbation

$$\begin{aligned}
\sum_{\mathbf{T}} e^{i\mathbf{q}\cdot\mathbf{T}} \frac{\partial \rho(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} &= 2 \sum_{\mathbf{T}n\mathbf{k}} e^{i\mathbf{q}\cdot\mathbf{T}} \left(\frac{\partial \varphi_{n\mathbf{k}}^*(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} \varphi_{n\mathbf{k}}(\mathbf{r}) + \varphi_{n\mathbf{k}}^*(\mathbf{r}) \frac{\partial \varphi_{n\mathbf{k}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} \right) \\
&= 2 \sum_{\mathbf{T}n\mathbf{k}} e^{i\mathbf{q}\cdot\mathbf{T}} \left(\frac{\partial \chi_{n\mathbf{k}}^*(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} \chi_{n\mathbf{k}}(\mathbf{r}) + \chi_{n\mathbf{k}}^*(\mathbf{r}) \frac{\partial \chi_{n\mathbf{k}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} \right) \\
&= 2e^{i\mathbf{q}\cdot\mathbf{r}} \sum_{\mathbf{T}n\mathbf{k}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r})} \left(\frac{\partial \chi_{n\mathbf{k}}^*(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} \chi_{n\mathbf{k}}(\mathbf{r}) + \chi_{n\mathbf{k}}^*(\mathbf{r}) \frac{\partial \chi_{n\mathbf{k}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} \right)
\end{aligned} \tag{39}$$

$$\begin{aligned}
&\left(-\frac{\nabla^2}{2} + V_{\text{KS}}(\mathbf{r}) - \varepsilon_{n\mathbf{k}} \right) \frac{\partial \varphi_{n\mathbf{k}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} = \left(\frac{\partial \varepsilon_{n\mathbf{k}}}{\partial R_{\mathbf{T}s\alpha}^0} - \frac{\partial V_{\text{KS}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} \right) \varphi_{n\mathbf{k}}(\mathbf{r}) \\
&\left(-\frac{(i\mathbf{k} + \nabla)^2}{2} + V_{\text{KS}}(\mathbf{r}) - \varepsilon_{n\mathbf{k}} \right) \frac{\partial \chi_{n\mathbf{k}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} = \left(\frac{\partial \varepsilon_{n\mathbf{k}}}{\partial R_{\mathbf{0}s\alpha}^0} - \frac{\partial V_{\text{KS}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} \right) \chi_{n\mathbf{k}}(\mathbf{r}) \\
\sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r})} \left(-\frac{(i\mathbf{k} + \nabla)^2}{2} + V_{\text{KS}}(\mathbf{r}) - \varepsilon_{n\mathbf{k}} \right) \frac{\partial \chi_{n\mathbf{k}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} &= \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r})} \left(\frac{\partial \varepsilon_{n\mathbf{k}}}{\partial R_{\mathbf{0}s\alpha}^0} - \frac{\partial V_{\text{KS}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} \right) \chi_{n\mathbf{k}}(\mathbf{r}) \\
\left(-\frac{(i\mathbf{k} + i\mathbf{q} + \nabla)^2}{2} + V_{\text{KS}}(\mathbf{r}) - \varepsilon_{n\mathbf{k}} \right) \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r})} \frac{\partial \chi_{n\mathbf{k}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} &= \left(N_C \delta_{\mathbf{q}\mathbf{0}} \frac{\partial \varepsilon_{n\mathbf{k}}}{\partial R_{\mathbf{0}s\alpha}^0} - \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r})} \frac{\partial V_{\text{KS}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} \right) \chi_{n\mathbf{k}}(\mathbf{r}).
\end{aligned} \tag{40}$$

Each component has lattice periodicity

$$\sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r}-\mathbf{T}')} \frac{\partial \chi_{n\mathbf{k}}(\mathbf{r} + \mathbf{T}')}{\partial R_{\mathbf{T}s\alpha}^0} = \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r})} \frac{\partial \chi_{n\mathbf{k}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} \tag{41}$$

$$\sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r}-\mathbf{T}')} \frac{\partial V_{\text{KS}}(\mathbf{r} + \mathbf{T}')}{\partial R_{\mathbf{T}s\alpha}^0} = \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r})} \frac{\partial V_{\text{KS}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} \tag{42}$$

Deformation potential is computed as follows:

$$\begin{aligned}
\sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r})} \frac{\partial V_{\text{KS}}(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} &= \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r})} \left\{ \frac{\partial V(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} + \int d^3r' \left(\frac{\delta V_{\text{H}}(\mathbf{r})}{\delta \rho(\mathbf{r}')} + \frac{\delta V_{\text{XC}}(\mathbf{r})}{\delta \rho(\mathbf{r}')} \right) \frac{\partial \rho(\mathbf{r}')}{\partial R_{\mathbf{T}s\alpha}^0} \right\} \\
&= \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r})} \left\{ \frac{\partial V(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} + \int d^3r' \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} + f_{\text{XC}}(\mathbf{r}, \mathbf{r}') \right) \frac{\partial \rho(\mathbf{r}')}{\partial R_{\mathbf{T}s\alpha}^0} \right\} \\
&= \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r})} \frac{\partial V(\mathbf{r})}{\partial R_{\mathbf{T}s\alpha}^0} + \int d^3r' e^{i\mathbf{q}\cdot(\mathbf{r}'-\mathbf{r})} \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} + f_{\text{XC}}(\mathbf{r}, \mathbf{r}') \right) \sum_{\mathbf{T}} e^{i\mathbf{q}\cdot(\mathbf{T}-\mathbf{r}')} \frac{\partial \rho(\mathbf{r}')}{\partial R_{\mathbf{T}s\alpha}^0}
\end{aligned} \tag{43}$$